

Dilution:

Disinfection 1:128 1 oz. per gallon of water
Sanitizer 1:256 1/2 oz. per gallon of water
SI USTED NO ENTIENDE LA ETIQUETA, BUSQUE A ALGUIEN PARA QUE SE LA EXPLIQUE A USTED EN DETALLE. If you do not understand the label, find someone to explain it to you in detail.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Preparation of Disinfectant Use-Solution: Add 1 oz. per gallon of water (1:128). Use 1 oz. per gallon of water to kill Norovirus, Enterobacter aerogenes.
DISINFECTION / VIRUCIDAL / FUNGICIDAL / MOLD AND MILDEW CONTROL
DIRECTIONS: Spray use-solution 6-8 inches from hard, nonporous surfaces, thoroughly wetting surfaces. Treated surfaces must remain visibly wet for 3 minutes. Do not breathe spray.

For disinfection against HIV-1 and Influenza Virus Type A / H3N2 and SARS-CoV-2, treated surfaces must remain visibly wet for 1 minute. For disinfection against Enterovirus and Norovirus, treated surfaces must remain visibly wet for 5 minutes. For disinfection against Legionella pneumophila and Adenovirus Type 5 and Adenovirus Type 7 and SARS Associated Coronavirus, treated surfaces must remain visibly wet for 10 minutes. Wipe dry with a cloth, sponge or mop or allow to air dry. For visibly soiled areas, a preliminary cleaning is required. Rinse all surfaces that come in contact with food such as countertops, appliances, tables and stovetops with potable water before reuse. Do not use on utensils, glassware and dishes.

For use as a One-Step Cleaner and Disinfectant by Electrostatic Spraying:
Electrostatic spray application is for bactericidal efficacy claims with a 3 minute contact time and 1:256 dilution and virucidal efficacy claims with a 5 minute contact time and 1:128 dilution. This application method is not to be used to treat surfaces for sanitization or fungicidal claims.

- Ensure bystanders and pets are not present in the room during application.
- Clean visibly dirty surfaces prior to spraying.
- Wear a N95 filtering facepiece respirator or half face respirator with a N95 filter and eye protection (goggles, face shield or safety glasses) when spraying.
- Select a spray nozzle that produces spray droplet particles with a volume median diameter (VMD) ≥ 80 µm.

- Place the electrostatic spray function in the ON position for electrostatic spray models that have the functionality to toggle ON/OFF.
- Hold the spray nozzle 12 inches away from the target surface. Starting at the highest point and working down to the lowest point, spray target surfaces using a smooth methodical "S" pattern. To avoid leaving gaps in sprayed surfaces, overlap previously treated surfaces when moving onto subsequent sections.
- The target surface must remain visibly wet for 5 minutes. Re-apply product as necessary to keep surface visibly wet for the duration of the contact time. Wipe surfaces dry or allow to air dry.

• Allow the treated room to remain unoccupied for 15 minutes before reentering.
Bactericidal Stability of Use-Dilution: Tests confirm that the use solution of this product, when stored in a sealed container such as a spray bottle, remains effective for 1 year. If product becomes visibly dirty or contaminated, the use-dilution must be discarded and fresh product prepared. Always use clean, properly labeled containers when diluting this product.

For Fungicidal Activity: This product is an effective fungicide against Trichophyton interdigitale (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities. This product is a one-step² fungicide at 1/2 oz. per gallon in 250 ppm hard water and 5% serum. **For Mold And Mildew:** At 1/2 oz. per gallon, this product will effectively inhibit the growth of mold and mildew and the odors caused by them when applied to hard, nonporous surfaces. Follow disinfection instructions. Repeat treatment every seven days, or more often if new growth appears.

CONTINUED FROM LEFT PANEL

This product is not to be used as a terminal sterilant / high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

TO SANITIZE SOFT SURFACES:

Add 1/2 - 2 oz. per gallon of water. Apply use-solution with a sprayer. Hold bottle 6" - 8" from surface and spray until fabric is wet. Fabric must remain visibly wet for 5 minutes. Allow fabric to Air dry. Reapply as necessary. Visibly soiled areas must be cleaned prior to sanitizing.
KILLS HIV AND HBV AND HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids, and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) associated with AIDS or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).
SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OR HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS. Personal Protection: Cleanup must always be done wearing protective gloves, gowns, masks and eye protection. **Cleaning Procedures:** Blood and other body fluids containing HIV-1 or HBV or HCV must be thoroughly cleaned from surfaces and objects before application of this product. **Contact Time:** Leave surface visibly wet for 1 minute for HIV-1 and 3 minutes for HBV and HCV with a 1/2 oz per gallon use-solution.

Use contact time indicated on the label for disinfection against all other viruses, bacteria and fungi claimed. **Disposal of Infectious Material:** Blood, body fluids, cleaning materials and clothing must be autoclaved and disposed of according to local regulations for infectious waste disposal.

TO SANITIZE NON-FOOD CONTACT SURFACES:

Spray use-solution 6-8 inches from hard, non-porous surfaces, visibly wetting surfaces. Treated surfaces must remain visibly wet for 15 seconds. Wipe dry with a sponge, mop or cloth or allow to air dry. Do not use in food preparation, food handling and food storage areas. This product is not to be used on glassware, dishware, cookware or utensils.

TO CLEAN AND DEODORIZE: Mix 1/2 oz. of this product per gallon of water to clean and deodorize hard, non-porous, non-food surfaces. Apply using a sprayer. Wipe or allow to air dry. Rinse all surfaces that come in contact with food such as countertops, exteriors of appliances, tables and stovetops with notatable water before reuse. Do not use on utensils, glassware and dishes.

DEODORING DIRECTIONS: Spray Solutions: 1/2-1 oz. per gallon of water eliminates undesirable odors. Allow surfaces to air dry. **Carpets:** 1/2-1 oz. per gallon of water eliminates odors associated with brine, vomit, smoke, and mildew. Reapply using a spray device after cleaning for freshening effects. Allow to dry.

Cleaning Solution: Dilute 1/2-1 oz. per gallon of water for cleaning of kitchen counters, tables, walls, bathrooms, toilet seats, floors and large routine applications. Apply to surfaces using trigger spray or other spray device. Remove excess liquid or allow surfaces to air dry.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
Pesticide Storage: Open dumping is prohibited. Store in original container in areas inaccessible to children. **Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **Container Disposal:** Non-refillable container. Do not reuse or refill this container. Wrap empty container and put in trash.

This product is a concentrated, Hospital Use disinfectant that is effective against a broad spectrum of bacteria, is virucidal* (including HIV-1, the AIDS Virus and Hepatitis B Virus and Hepatitis C Virus), fungicidal, and inhibits the growth of mold and mildew, and their odors, when used as directed. Effective against SARS-Related Coronavirus 2* SARS-CoV-2 in 1 minute. Kills HIV virus in just 1 minute contact time. Kills flu virus in just 1 minute contact time. Kills Rotavirus in 3 minutes. Leaves a clean fresh scent. Tested according to ADAC Germicidal Spray Products as Disinfectants test. Cross-contamination is of major housekeeping concern. This product has been formulated to aid in the reduction of cross-contamination on treated surfaces not only in hospitals, but in schools, institutions and industry. This product Sanitizes² 99.9% of Staphylococcus aureus and Klebsiella pneumoniae in 15 seconds.

For sanitizing and disinfecting of ultrasound transducers, probes, mammography compressor plates and other hard nonporous surfaces. To clean and disinfect hard, nonporous finished floors. Cleans and disinfects without dulling gloss.

AREAS OF USE: Use this product in: clinics, dental offices, health care facilities, hospitals, medical offices, medical related facilities, nursing homes and sick rooms. Use in business and office buildings, colleges, institutional facilities, public facilities, ballrooms, restrooms, public rest rooms, schools, universities, ambulances and emergency EMS vehicles.

FOR MEDICAL EMERGENCIES, CALL

1-888-779-1339.

FOR CUSTOMER SERVICE

INFORMATION 1-800-323-2251

www.triple-s.com

Product No.
13215



TRIPLE S

NAVIGATOR
Dilution Control System

3x
RENEGADE DAILY ONE-STEP DISINFECTANT

Staphylocidal⁶ • Pseudomonacidal⁷ • Bactericidal • Fungicidal • Virucidal*

Kills SARS-CoV-2* virus, the cause of COVID-19

Active Ingredients:
Octyl decyl dimethyl ammonium chloride.....6.67%
Dioctyl dimethyl ammonium chloride.....2.67%
Didecyl dimethyl ammonium chloride.....4.00%
Alkyl (C14, 50%; C12, 40%; C16, 10%) dimethyl benzy ammonium chloride.....8.90%
Other Ingredients:.....77.76%
Total.....100.00%

KEEP OUT OF REACH OF CHILDREN DANGER PELIGRO
See back panel for Precautionary Statements.
E.P.A. Reg. No. 6836-349-12120
E.P.A. Est. No. 8325-PA-01

FIRST AID
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling poison control center or doctor or going for treatment. For general information on product use, etc., call the National Pesticides Information Center at 1-800-858-7378. You may also contact the poison control center at 1-800-222-1222 for emergency medical treatment information.

NET CONTENTS: 67.8 fl. oz. (2 Liters)



TRIPLE S

NAVIGATOR
Dilution Control System

3x
RENEGADE DAILY ONE-STEP DISINFECTANT

Staphylocidal⁶ • Pseudomonacidal⁷ • Bactericidal • Fungicidal • Virucidal*

Kills SARS-CoV-2* virus, the cause of COVID-19



TRIPLE S