

OXYPRO 365

HYDROGEN PEROXIDE DISINFECTANT CLEANER

Technical Data

Areas of Use

- Countertops
- Bathroom fixtures
- Tables
- Desks

Specifications

SSS Item Numbers	48411, 48412
Available Sizes	1Qt & 1G
Color	Clear
Fragrance	Fresh
Physical State	Liquid
Base	Hydrogen peroxide
pH	1.5
Viscosity	Water thin
Foam	Moderate
Abrasives	None
Solubility	Complete with water
Recommended Dilution	RTU
Fixed Dilution	RTU
Fungicidal	Yes
Detergency	Excellent
Rinsability	Excellent
Flash Point	None (waterbased)
Hard Water Tolerance	Good
Biodegradability	Complete
Storage/Stability	1 year
Weight Per Gallon	8.46 lbs./gal.
Freeze/Thaw Stability	Keep from freezing

HMIS Hazardous Materials I.D. System

Health	0
Flammability	0
Reactivity	0
Personal Protection	-

What the Pros Know

Effectively cleans and deodorizes surfaces in restrooms, and other areas where bacterial growth can cause malodors.

Description

This product is a one-step RTU hydrogen peroxide disinfectant cleaner that is effective against a broad spectrum of bacteria, is virucidal* (including HIV-1, the AIDS Virus, Methicillin Resistant Staphylococcus aureus (MSRA), in 3 minutes. Kills 99.9% SARS-CoV-2 (COVID-19) in 1 minute. This product is bleach and chlorine free. Leaves no harsh or sticky residue. This product sanitizes in 1 minute against Staphylococcus aureus and Enterobacter aerogenes on hard, non-porous, non-food contact surfaces. Inhibits the growth of mold and mildew, leaving a clean and fresh fragrance. Requires no PPE.

Directions

GENERAL CLEANING / DEODORIZING DIRECTIONS: Apply to soiled area then wipe clean. Repeat for visibly soiled areas.

DISINFECTION/*VIRUCIDAL DIRECTIONS: Spray 6-8 inches from surface, until surfaces are visibly wet. Do not breathe spray. Treated surfaces must remain visibly wet for 3 minutes. For SARS-CoV-2, treated surfaces must remain visibly wet for 1 minute. Wipe dry with a clean cloth, sponge, mop or towel, or allow to air dry.

MOLD AND MILDEW CONTROL DIRECTIONS: This product effectively inhibits the growth of mold and mildew, and the odors caused by them, when applied to hard, non-porous surfaces. Spray 6 - 8 inches from surface, until surfaces are visibly wet. Do not breathe spray. Allow to air dry on surface. Repeat every 7 days, or when mold or mildew growth returns.

This product is not for use on medical device surfaces.

NON-ACID TOILET BOWL AND URINAL CLEANER DIRECTIONS:

Apply this product to exposed surfaces including under the rim with a sprayer. Use a toilet brush to scrub the bowl. Repeat and allow to remain visibly wet for 1 minute, then flush.

***KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings, or other settings, in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES / OBJECTS SOILED WITH BLOOD / BODY FLUIDS:

PERSONAL PROTECTION: Clean-up must always be done wearing protective gloves, gowns, masks and eye protection.

CLEANING PROCEDURES: Blood and other body fluids containing HIV must be thoroughly cleaned from surfaces and objects before application of this product. **CONTACT TIME:** Leave surface visibly wet for 3 minutes.

DISPOSAL OF INFECTIOUS MATERIALS: Blood, body fluids, cleaning materials and clothing must be autoclaved and disposed of according to local regulations for infectious waste disposal.

TO SANITIZE AND DEODORIZE NON-FOOD CONTACT SURFACES:

Spray this product until visibly wet. Treated hard, non-porous, non-food surfaces must remain visibly wet for 1 minute. Wipe dry with a clean cloth, sponge, mop or towel, or allow to air dry. For visibly soiled areas, pre-cleaning is required.

*Cleans, Deodorizes
and Disinfects
in one-step*

*Broad range of
kill claims*

*Economical to
Use*

*Safe to Use and
Handle*

